

Three dimensional pseudomanifolds on eight vertices

Basudeb Datta,¹ and Nandini Nilakantan²

¹Department of Mathematics, Indian Institute of Science, Bangalore 560 012, India
dattab@math.iisc.ernet.in

²Department of Mathematics & Statistics, Indian Institute of Technology, Kanpur 208 016, India
nandini@iitk.ac.in

To appear in ‘International Journal of Mathematics and Mathematical Sciences’

Abstract. A normal pseudomanifold is a pseudomanifold in which the links of simplices are also pseudomanifolds. So, a normal 2-pseudomanifold triangulates a connected closed 2-manifold. But, normal d -pseudomanifolds form a broader class than triangulations of connected closed d -manifolds for $d \geq 3$. Here, we classify all the 8-vertex neighbourly normal 3-pseudomanifolds. This gives a classification of all the 8-vertex normal 3-pseudomanifolds. There are 73 such 3-pseudomanifolds, 38 of which triangulate the 3-sphere and other 35 are not combinatorial 3-manifolds. These 35 triangulate six distinct topological spaces. As a preliminary result, we show that any 8-vertex 3-pseudomanifold is equivalent by proper bistellar moves to an 8-vertex neighbourly 3-pseudomanifold. This result is the best possible since there exists a 9-vertex non-neighbourly 3-pseudomanifold (B_9^3 in Example 7 below) which does not allow any proper bistellar moves.

Mathematics Subject Classification (2000): 57Q15, 57Q05, 57N05, 55M25.

Keywords: Bistellar moves; Normal pseudomanifold; Triangulated 3-sphere; Singular vertex.

1 Introduction

Recall that a *simplicial complex* is a collection of non-empty finite sets (sets of *vertices*) such that every non-empty subset of an element is also an element. For $i \geq 0$, the elements of size $i + 1$ are called the *i -simplices* (or *i -faces*) of the complex.

A simplicial complex is usually thought of as a prescription for construction of a topological space by pasting geometric simplices. The space thus obtained from a simplicial complex K is called the *geometric carrier* of K and is denoted by $|K|$. We also say that K *triangulates* $|K|$. A *combinatorial 2-manifold* (respectively, *combinatorial 2-sphere*) is a simplicial complex which triangulates a closed surface (respectively, the 2-sphere S^2).

For a simplicial complex K , the maximum of k such that K has a k -simplex is called the *dimension* of K . A d -dimensional simplicial complex K is called *pure* if each simplex of K is contained in a d -simplex of K . A d -simplex in a pure d -dimensional simplicial complex is called a *facet*. A d -dimensional pure simplicial complex K is called a *weak pseudomanifold* if each $(d - 1)$ -simplex of K is contained in exactly two facets of K .

With a pure simplicial complex K of dimension $d \geq 1$, we associate a graph $\Lambda(K)$ as follows. The vertices of $\Lambda(K)$ are the facets of K and two vertices of $\Lambda(K)$ are adjacent if the corresponding facets intersect in a $(d - 1)$ -simplex of K . If $\Lambda(K)$ is connected then K is called *strongly connected*. A strongly connected weak pseudomanifold is called a *pseudomanifold*. Thus, for a d -pseudomanifold K , $\Lambda(K)$ is a connected $(d + 1)$ -regular graph. This implies

that K has no proper subcomplex which is also a d -pseudomanifold. (Or else, the facets of such a subcomplex would provide a disconnection of $\Lambda(X)$.)

For any set V with $\#(V) = d + 2$ ($d \geq 0$), let K be the simplicial complex whose simplexes are all the non-empty proper subsets of V . Then K is a d -pseudomanifold and triangulates the d -sphere S^d . This d -pseudomanifold K is called the *standard d -sphere* and is denoted by $S_{d+2}^d(V)$ (or S_{d+2}^d). By convention, S_2^0 is the only 0-pseudomanifold.

If σ is a face of a simplicial complex K then the *link* of σ in K , denoted by $\text{lk}_K(\sigma)$ (or $\text{lk}(\sigma)$), is by definition the simplicial complex whose faces are the faces τ of K such that τ is disjoint from σ and $\sigma \cup \tau$ is a face of K . Clearly, the link of an i -face in a weak d -pseudomanifold is a weak $(d - i - 1)$ -pseudomanifold. For $d \geq 1$, a connected weak d -pseudomanifold is said to be a *normal d -pseudomanifold* if the links of all the simplices of dimension $\leq d - 2$ are connected. Thus, any connected triangulated d -manifold (triangulation of a closed d -manifold) is a normal d -pseudomanifold. Clearly, the normal 2-pseudomanifolds are just the connected combinatorial 2-manifolds. But, normal d -pseudomanifolds form a broader class than connected triangulated d -manifolds for $d \geq 3$.

Observe that if X is a normal pseudomanifold then X is a pseudomanifold. (If $\Lambda(X)$ is not connected then, since X is connected, $\Lambda(X)$ has two components G_1 and G_2 and two intersecting facets σ_1, σ_2 such that $\sigma_i \in G_i$, $i = 1, 2$. Choose σ_1, σ_2 among all such pairs such that $\dim(\sigma_1 \cap \sigma_2)$ is maximum. Then $\dim(\sigma_1 \cap \sigma_2) \leq d - 2$ and $\text{lk}_X(\sigma_1 \cap \sigma_2)$ is not connected, a contradiction.) Notice that all the links of positive dimensions (i.e., the links of simplices of dimension $\leq d - 2$) in a normal d -pseudomanifold are normal pseudomanifolds. Thus, if K is a normal 3-pseudomanifold then the link of a vertex in K is a combinatorial 2-manifold. A vertex v of a normal 3-pseudomanifold K is called *singular* if the link of v in K is not a 2-sphere. The set of singular vertices is denoted by $\text{SV}(K)$. Clearly, the space $|K| \setminus \text{SV}(K)$ is a pl 3-manifold. If $\text{SV}(K) = \emptyset$ (i.e., the link of each vertex is a 2-sphere) then K is called a *combinatorial 3-manifold*. A *combinatorial 3-sphere* is a combinatorial 3-manifold which triangulates the topological 3-sphere S^3 .

Let M be a weak d -pseudomanifold. If α is a $(d - i)$ -face of M , $0 < i \leq d$, such that $\text{lk}_M(\alpha) = S_{i+1}^{i-1}(\beta)$ and β is not a face of M (such a face α is said to be a *removable face* of M) then consider the weak d -pseudomanifold (denoted by $\kappa_\alpha(M)$) whose facet-set is $\{\sigma : \sigma \text{ a facet of } M, \alpha \not\subseteq \sigma\} \cup \{\beta \cup \alpha \setminus \{v\} : v \in \alpha\}$. The operation $\kappa_\alpha : M \mapsto \kappa_\alpha(M)$ is called a *bistellar i -move*. For $0 < i < d$, a bistellar i -move is called a *proper bistellar move*. If κ_α is a proper bistellar i -move and $\text{lk}_M(\alpha) = S_{i+1}^{i-1}(\beta)$ then β is a removable i -face of $\kappa_\alpha(M)$ (with $\text{lk}_{\kappa_\alpha(M)}(\beta) = S_{d-i+1}^{d-i-1}(\alpha)$) and $\kappa_\beta : \kappa_\alpha(M) \mapsto M$ is a bistellar $(d - i)$ -move. For a vertex u , if $\text{lk}_M(u) = S_{d+1}^{d-1}(\beta)$ then the bistellar d -move $\kappa_{\{u\}} : M \mapsto \kappa_{\{u\}}(M) = N$ deletes the vertex u (we also say that N is obtained from M by *collapsing* the vertex u). The operation $\kappa_\beta : N \mapsto M$ is called a *bistellar 0-move* (we also say that M is obtained from N by *starring* the vertex u in the facet β of N). The 10-vertex combinatorial 3-manifold A_{10}^3 in Example 6 is not neighbourly and does not allow any bistellar 1-move. In [5], Bagchi and Datta have shown that if the number of vertices in a non-neighbourly combinatorial 3-manifold is at most 9 then the 3-manifold admits a bistellar 1-move. Existence of the 9-vertex 3-pseudomanifold B_9^3 in Example 7 shows that Bagchi and Datta's result is not true for 9-vertex 3-pseudomanifolds. Here we prove.

Theorem 1. *If M is an 8-vertex 3-pseudomanifold, then there exist a sequence of bistellar 1-moves $\kappa_{A_1}, \dots, \kappa_{A_m}$, for some $m \geq 0$, such that $\kappa_{A_m}(\dots(\kappa_{A_1}(M)))$ is a neighbourly 3-pseudomanifold.*

In [1], Altshuler has shown that every combinatorial 3-manifold with at most 8 vertices

is a combinatorial 3-sphere. In [10], Grünbaum and Sreedharan have shown that there are exactly 37 polytopal 3-spheres on 8 vertices (namely, $S_{8,1}^3, \dots, S_{8,37}^3$ in Examples 2 and 3). They have also constructed the non-polytopal sphere $S_{8,38}^3$. In [6], Barnette proved that there is only one more non-polytopal 8-vertex 3-sphere (namely, $S_{8,39}^3$). In [8], Emch constructed a 8-vertex normal 3-pseudomanifold (namely, N_1 in Example 4) as a block design. This is not a combinatorial 3-manifold and its automorphism group is $\text{PGL}(2, 7)$ (cf. [13]). In [2], Altshuler has constructed another 8-vertex normal 3-pseudomanifold (namely, N_5 in Example 4). In [14], Lutz has shown that there exist exactly three 8-vertex normal 3-pseudomanifolds which are not combinatorial 3-manifolds (namely, N_1 , N_5 and N_6 in Example 4) with vertex-transitive automorphism groups. Here we prove:

Theorem 2. *Let $S_{8,35}^3, \dots, S_{8,38}^3, N_1, \dots, N_{15}$ be as in Examples 2 and 4.*

- (a) *Then $S_{8,i}^3 \not\cong S_{8,j}^3$, $N_k \not\cong N_l$ and $S_{8,m}^3 \not\cong N_n$ for $35 \leq i < j \leq 38$, $1 \leq k < l \leq 15$, $35 \leq m \leq 38$ and $1 \leq n \leq 15$.*
- (b) *If M is an 8-vertex neighbourly normal 3-pseudomanifold then M is isomorphic to one of $S_{8,35}^3, \dots, S_{8,38}^3, N_1, \dots, N_{15}$.*

Corollary 3. *There are exactly 39 combinatorial 3-manifolds on 8 vertices, all of which are combinatorial 3-spheres.*

Corollary 4. *There are exactly 35 normal 3-pseudomanifolds on 8 vertices which are not combinatorial 3-manifolds. These are $N_1, \dots, N_{35}$ defined in Examples 4 and 5.*

The topological properties of these normal 3-pseudomanifolds are given in Section 3.

2 Preliminaries

All the simplicial complexes considered in this paper are finite (i.e., with finite vertex-set). The vertex-set of a simplicial complex K is denoted by $V(K)$. We identify the 0-faces of a complex with the vertices. The 1-faces of a complex K are also called the *edges* of K .

If K, L are two simplicial complexes, then an *isomorphism* from K to L is a bijection $\pi : V(K) \rightarrow V(L)$ such that for $\sigma \subseteq V(K)$, σ is a face of K if and only if $\pi(\sigma)$ is a face of L . Two complexes K, L are called *isomorphic* when such an isomorphism exists. We identify two complexes if they are isomorphic. An isomorphism from a complex K to itself is called an *automorphism* of K . All the automorphisms of K form a group under composition, which is denoted by $\text{Aut}(K)$.

For a face σ in a simplicial complex K , the number of vertices in $\text{lk}_K(\sigma)$ is called the *degree* of σ in K and is denoted by $\deg_K(\sigma)$ (or by $\deg(\sigma)$). If every pair of vertices of a simplicial complex K form an edge then K is called *neighbourly*. For a simplicial complex K , if $U \subseteq V(K)$ then $K[U]$ denotes the induced complex of K on the vertex-set U .


If the number of i -faces of a d -dimensional simplicial complex K is $f_i(K)$ ($0 \leq i \leq d$), then the number $\chi(K) := \sum_{i=0}^d (-1)^i f_i(K)$ is called the *Euler characteristic* of K .

A *graph* is a simplicial complex of dimension ≤ 1 . A finite 1-pseudomanifold is called a *cycle*. An n -cycle is a cycle on n vertices and is denoted by C_n (or by $C_n(a_1, \dots, a_n)$ if the edges are $a_1a_2, \dots, a_{n-1}a_n, a_na_1$).

For a simplicial complex K , the graph consisting of the edges and vertices of K is called the *edge-graph* of K and is denoted by $\text{EG}(K)$. The complement of $\text{EG}(K)$ is called the

non-edge graph of K and is denoted by $\text{NEG}(K)$. For a weak 3-pseudomanifold M and an integer $n \geq 3$, we define the graph $G_n(M)$ as follows. The vertices of $G_n(M)$ are the vertices of M . Two vertices u and v form an edge in $G_n(M)$ if uv is an edge of degree n in M . Clearly, if M and N are isomorphic then $G_n(M)$ and $G_n(N)$ are isomorphic for each n .

If M is a weak 3-pseudomanifold and $\kappa_\alpha : M \mapsto \kappa_\alpha(M) = N$ is a bistellar 1-move then, from the definition, $(f_0(N), f_1(N), f_2(N), f_3(N)) = (f_0(M), f_1(M)+1, f_2(M)+2, f_3(M)+1)$ and $\deg_N(v) \geq \deg_M(v)$ for any vertex v . If $\kappa_\alpha : M \mapsto \kappa_\alpha(M) = L$ is a bistellar 3-move then $(f_0(L), f_1(L), f_2(L), f_3(L)) = (f_0(M) - 1, f_1(M) - 4, f_2(M) - 6, f_3(M) - 3)$.



Bistellar moves in dimension 3

Consider the binary relation ' $\leq$ ' on the set of weak 3-pseudomanifolds as: $M \leq N$ if there exists a finite sequence of bistellar 1-moves $\kappa_{\alpha_1}, \dots, \kappa_{\alpha_m}$, for some $m \geq 0$, such that $N = \kappa_{\alpha_m}(\dots \kappa_{\alpha_1}(M))$. Clearly, this $\leq$ is a partial order relation.

Two weak d -pseudomanifolds M and N are *bistellar equivalent* (denoted by $M \sim N$) if there exists a finite sequence of bistellar operations leading from M to N . If there exists a finite sequence of proper bistellar operations leading from M to N then we say M and N are *properly bistellar equivalent* and we denote this by $M \approx N$. Clearly, ' $\sim$ ' and ' $\approx$ ' are equivalence relations on the set of pseudomanifolds. It is easy to see that $M \sim N$ implies that $|M|$ and $|N|$ are pl homeomorphic.

For two simplicial complexes X and Y with disjoint vertex sets, the simplicial complex $X * Y := X \cup Y \cup \{\sigma \cup \tau : \sigma \in X, \tau \in Y\}$ is called the *join* of X and Y .

Let K be an n -vertex (weak) d -pseudomanifold. If u is a vertex of K and v is not a vertex of K then consider the simplicial complex $\Sigma_{uv}K$ on the vertex set $V(K) \cup \{v\}$ whose set of facets is $\{\sigma \cup \{u\} : \sigma \text{ is a facet of } K \text{ and } u \notin \sigma\} \cup \{\tau \cup \{v\} : \tau \text{ is a facet of } K\}$. Then $\Sigma_{uv}K$ is a (weak) $(d+1)$ -pseudomanifold and $|\Sigma_{uv}K|$ is the topological suspension $S|K|$ of $|K|$ (cf. [3]). Easy to see that the links of u and v in $\Sigma_{uv}K$ are isomorphic to K . This $\Sigma_{uv}K$ is called the *one-point suspension* of K .

For two d -pseudomanifolds X and Y , a simplicial map $f: X \rightarrow Y$ is called a *k-fold branched covering* (with discrete branch locus) if $|f|_{|X| \setminus f^{-1}(U)}: |X| \setminus f^{-1}(U) \rightarrow |Y| \setminus U$ is a k -fold covering for some $U \subseteq V(Y)$. (We say that X is a *branched cover* of Y and Y is a *branched quotient* of X .) The smallest such U (so that $|f|_{|X| \setminus f^{-1}(U)}: |X| \setminus f^{-1}(U) \rightarrow |Y| \setminus U$ is a covering) is called the *branch locus*. If N is a k -fold branched quotient of M and $\tilde{N}$ is obtained from N by collapsing a vertex (respectively starring a vertex in a facet) then $\tilde{N}$ is the branched quotient of $\tilde{M}$, where $\tilde{M}$ can be obtained from M by collapsing k vertices (respectively starring k vertices in k facets). For proper bistellar moves we have:


Lemma 2.1. *Let M and N be two d -pseudomanifolds and $f: M \rightarrow N$ be a k -fold branched covering. For $1 \leq l < d-1$, if α is a removable l -face, then $f^{-1}(\alpha)$ consists of k removable l -faces $\alpha_1, \dots, \alpha_k$ (say) and $\kappa_{\alpha_k}(\dots(\kappa_{\alpha_1}(M)))$ is a k -fold branched cover of $\kappa_\alpha(N)$.*

Proof. Let $\text{lk}_N(\alpha) = S_{d-l+1}^{d-l-1}(\beta)$. Since the dimension of α is > 0 , $f^{-1}(\alpha)$ consists of k l -faces, $\alpha_1, \dots, \alpha_k$ (say) of M . Let $\text{lk}_M(\alpha_i) = S_{d-l+1}^{d-l-1}(\beta_i)$ and $M_i := M[\alpha_i \cup \beta_i]$ for $1 \leq i \leq k$. Since f is simplicial, β_i is not a face of M and hence α_i is removable for each i . Since

$0 < l < d - 1$, it follows that M_i is neighbourly. For $i \neq j$, if $x \neq y \in V(M_i) \cap V(M_j)$, then xy is an edge in $M_i \cap M_j$ and hence the number of edges in $f^{-1}(f(x)f(y))$ is less than k , a contradiction. So, $\#(V(M_i) \cap V(M_j)) \leq 1$ for $i \neq j$. This implies that β_i is not a face in $\kappa_{\alpha_j}(M)$ and hence α_i is removable in $\kappa_{\alpha_j}(M)$ for $i \neq j$. The result now follows. $\square$

Remark 1 shows that Lemma 2.1 is not true for $l = d - 1$ (i.e., for bistellar 1-moves) in general.

Example 1: Some weak 2-pseudomanifolds on at most seven vertices. The degree sequences are presented parenthetically below the figures. Each of $S_1, \dots, S_9$ triangulates the 2-sphere, each of $R_1, \dots, R_4$ triangulates the real projective plane and T triangulates the torus. Observe that P_1, P_2 are not pseudomanifolds.



We know that if K is a weak 2-pseudomanifold with at most six vertices then K is isomorphic to $S_1, \dots, S_4$ or R_1 (cf. [3]). In [7], we have seen the following.

Proposition 2.2. *There are exactly 13 distinct 2-dimensional weak pseudomanifolds on 7 vertices, namely, $S_5, \dots, S_9$, $R_2, \dots, R_4$, T , $P_1, \dots, P_3$ and P_4 .*

3 Examples

We identify a weak pseudomanifold with the set of facets in it.

Example 2: These four neighbourly 8-vertex combinatorial 3-manifolds were found by Grünbaum and Sreedharan (in [10], these are denoted by P_{35}^8 , P_{36}^8 , P_{37}^8 and $\mathcal{M}$ respectively). It follows from Lemma 3.2 that these are combinatorial 3-spheres. It was shown in [10] that the first three of these are polytopal 3-spheres and the last one is a non-polytopal sphere.

$$\begin{aligned}
S_{8,35}^3 &= \{1234, 1267, 1256, 1245, 2345, 2356, 2367, 3467, 3456, 4567, \\
&\quad 1238, 1278, 2378, 1348, 3478, 1458, 4578, 1568, 1678, 5678\}, \\
S_{8,36}^3 &= \{1234, 1256, 1245, 1567, 2345, 2356, 2367, 3467, 3456, 4567, \\
&\quad 1268, 1678, 2678, 1238, 2378, 1348, 3478, 1458, 1578, 4578\}, \\
S_{8,37}^3 &= \{1234, 1256, 1245, 1457, 2345, 2356, 2367, 3467, 3456, 4567, \\
&\quad 1568, 1578, 5678, 1268, 2678, 1238, 2378, 1348, 1478, 3478\}, \\
S_{8,38}^3 &= \{1234, 1237, 1267, 1347, 1567, 2345, 2367, 3467, 3456, 4567, \\
&\quad 2358, 2368, 3568, 1268, 1568, 1248, 2458, 1478, 1578, 4578\}.
\end{aligned}$$

Lemma 3.1. $S_{8,i}^3 \not\cong S_{8,j}^3$ for $35 \leq i < j \leq 38$.

Proof. Observe that $G_6(S_{8,35}^3) = C_8(1, 2, \dots, 8)$, $G_6(S_{8,36}^3) = (V, \{23, 34, 45, 67, 78, 81\})$, $G_6(S_{8,37}^3) = (V, \{23, 34, 56, 78, 81\})$ and $G_6(S_{8,38}^3) = (V, \{17, 23, 58\})$, where $V = \{1, \dots, 8\}$. Since $K \cong L$ implies $G_6(K) \cong G_6(L)$, $S_{8,i}^3 \not\cong S_{8,j}^3$, for $35 \leq i < j \leq 38$. $\square$

Example 3: Some non-neighbourly 8-vertex combinatorial 3-manifolds. It follows from Lemma 3.2 that these are combinatorial 3-spheres. For $1 \leq i \leq 34$, the sphere $S_{8,i}^3$ is isomorphic to the polytopal sphere P_i^8 in [10] and the sphere $S_{8,39}^3$ is isomorphic to the non-polytopal sphere found by Barnette in [6]. We consecutively define:

$$\begin{aligned}
S_{8,39}^3 &= \kappa_{46}(S_{8,38}^3), & S_{8,33}^3 &= \kappa_{27}(S_{8,37}^3), & S_{8,32}^3 &= \kappa_{48}(S_{8,37}^3), & S_{8,31}^3 &= \kappa_{58}(S_{8,37}^3), \\
S_{8,30}^3 &= \kappa_{24}(S_{8,37}^3), & S_{8,29}^3 &= \kappa_{27}(S_{8,31}^3), & S_{8,28}^3 &= \kappa_{24}(S_{8,31}^3), & S_{8,27}^3 &= \kappa_{13}(S_{8,31}^3), \\
S_{8,25}^3 &= \kappa_{57}(S_{8,31}^3), & S_{8,24}^3 &= \kappa_{48}(S_{8,31}^3), & S_{8,23}^3 &= \kappa_{35}(S_{8,31}^3), \\
S_{8,26}^3 &= \kappa_{46}(S_{8,27}^3), & S_{8,22}^3 &= \kappa_{24}(S_{8,25}^3), & S_{8,21}^3 &= \kappa_{68}(S_{8,25}^3), & S_{8,20}^3 &= \kappa_{48}(S_{8,25}^3), \\
S_{8,19}^3 &= \kappa_{17}(S_{8,25}^3), & S_{8,18}^3 &= \kappa_{27}(S_{8,25}^3), & S_{8,12}^3 &= \kappa_{15}(S_{8,25}^3), & S_{8,11}^3 &= \kappa_{35}(S_{8,25}^3), \\
S_{8,17}^3 &= \kappa_{24}(S_{8,19}^3), & S_{8,34}^3 &= \kappa_{27}(S_{8,26}^3) = S_3^0(1, 3) * S_3^0(2, 7) * S_3^0(4, 6) * S_3^0(5, 8), \\
S_{8,16}^3 &= \kappa_{13}(S_{8,19}^3), & S_{8,15}^3 &= \kappa_{28}(S_{8,18}^3), & S_{8,14}^3 &= \kappa_{47}(S_{8,20}^3), & S_{8,10}^3 &= \kappa_{15}(S_{8,19}^3), \\
S_{8,9}^3 &= \kappa_{35}(S_{8,19}^3), & S_{8,8}^3 &= \kappa_{47}(S_{8,19}^3), & S_{8,13}^3 &= \kappa_{38}(S_{8,16}^3), & S_{8,7}^3 &= \kappa_{24}(S_{8,8}^3), \\
S_{8,6}^3 &= \kappa_{35}(S_{8,8}^3), & S_{8,5}^3 &= \kappa_{48}(S_{8,8}^3), & S_{8,4}^3 &= \kappa_{15}(S_{8,8}^3), \\
S_{8,3}^3 &= \kappa_{48}(S_{8,4}^3), & S_{8,2}^3 &= \kappa_{48}(S_{8,6}^3), & S_{8,1}^3 &= \kappa_{16}(S_{8,4}^3).
\end{aligned}$$

Lemma 3.2. (a) $S_{8,i}^3 \approx S_{8,j}^3$, for $1 \leq i, j \leq 39$, (b) $S_{8,m}^3$ is a combinatorial 3-sphere, for $1 \leq m \leq 39$ and (c) $S_{8,k}^3 \not\cong S_{8,l}^3$, for $1 \leq k < l \leq 39$.

Proof. For $0 \leq i \leq 6$, let $\mathcal{S}_i$ denote the set of $S_{8,j}^3$'s with i non-edges. Then $\mathcal{S}_0 = \{S_{8,35}^3, S_{8,36}^3, S_{8,37}^3, S_{8,38}^3\}$, $\mathcal{S}_1 = \{S_{8,30}^3, S_{8,31}^3, S_{8,32}^3, S_{8,33}^3, S_{8,39}^3\}$, $\mathcal{S}_2 = \{S_{8,23}^3, S_{8,24}^3, S_{8,25}^3, S_{8,27}^3, S_{8,28}^3, S_{8,29}^3\}$, $\mathcal{S}_3 = \{S_{8,11}^3, S_{8,12}^3, S_{8,18}^3, S_{8,19}^3, S_{8,20}^3, S_{8,21}^3, S_{8,22}^3, S_{8,26}^3\}$, $\mathcal{S}_4 = \{S_{8,8}^3, S_{8,9}^3, S_{8,10}^3, S_{8,14}^3, S_{8,15}^3, S_{8,16}^3, S_{8,17}^3, S_{8,34}^3\}$, $\mathcal{S}_5 = \{S_{8,4}^3, S_{8,5}^3, S_{8,6}^3, S_{8,7}^3, S_{8,13}^3\}$ and $\mathcal{S}_6 = \{S_{8,1}^3, S_{8,2}^3, S_{8,3}^3\}$.

From the proof of Lemma 4.7, $S_{8,35}^3 \approx S_{8,30}^3 \approx S_{8,36}^3 \approx S_{8,30}^3 \approx S_{8,37}^3 \approx S_{8,32}^3 \approx S_{8,38}^3$. Thus, $S_{8,i}^3 \approx S_{8,j}^3$ for $35 \leq i, j \leq 38$. Now, if $S_{8,i}^3 \in \mathcal{S}_2 \cup \mathcal{S}_3 \cup \mathcal{S}_4 \cup \mathcal{S}_5 \cup \mathcal{S}_6$ then, from the definition of $S_{8,i}^3$, $S_{8,i}^3 \approx S_{8,31}^3 \approx S_{8,37}^3$. This proves Part (a).

Since $S_{8,34}^3$ is a join of spheres, $S_{8,34}^3$ is a combinatorial 3-sphere. Clearly, if $M \approx N$ and M is a combinatorial 3-sphere then N is so. Part (b) now follows from Part (a).

Since the non-edge graphs of the members of $\mathcal{S}_6$ (respectively, $\mathcal{S}_5$) are pairwise non-isomorphic, the members of $\mathcal{S}_6$ (respectively, $\mathcal{S}_5$) are pairwise non-isomorphic.

For $S_{8,i}^3, S_{8,j}^3 \in \mathcal{S}_4$ ($i < j$) and $\text{NEG}(S_{8,i}^3) \cong \text{NEG}(S_{8,j}^3)$ imply $(i, j) = (8, 9)$ or $(14, 15)$. Since $M \cong N$ implies $G_6(M) \cong G_6(N)$ and $G_6(S_{8,8}^3) \not\cong G_6(S_{8,9}^3)$, $G_6(S_{8,14}^3) \not\cong G_6(S_{8,15}^3)$, the members of $\mathcal{S}_4$ are pairwise non-isomorphic.

For $S_{8,i}^3 \neq S_{8,j}^3 \in \mathcal{S}_3$ and $\text{NEG}(S_{8,i}^3) \cong \text{NEG}(S_{8,j}^3)$ imply $\{i, j\} = \{11, 12\}$ or $18 \leq i \neq j \leq 21$. Since the non-edge graph of a member in Σ_i is non-isomorphic to the non-edge graph of a member of Σ_j for $i \neq j$, a member of Σ_i is non-isomorphic to a member of Σ_j . Observe that $G_6(S_{8,11}^3) \not\cong G_6(S_{8,12}^3)$ and for $18 \leq i < j \leq 21$, $G_6(S_{8,i}^3) \cong G_6(S_{8,j}^3)$ implies $(i, j) = (18, 19)$. Since $G_3(S_{8,18}^3) \not\cong G_3(S_{8,19}^3)$ the members of $\mathcal{S}_3$ are pairwise non-isomorphic.

Since $G_3(S_{8,i}^3) \not\cong G_3(S_{8,j}^3)$ for $S_{8,i}^3 \neq S_{8,j}^3 \in \mathcal{S}_2$, the members of $\mathcal{S}_2$ are pairwise non-isomorphic. By the same reasoning, the members of $\mathcal{S}_1$ are pairwise non-isomorphic.

By Lemma 3.1, the members of $\mathcal{S}_0$ are pairwise non-isomorphic. Since a member of $\mathcal{S}_i$ is non-isomorphic to a member of $\mathcal{S}_j$ for $i \neq j$, the above imply Part (c). $\square$

Example 4: Some 8-vertex neighbourly normal 3-pseudomanifolds.

$$\begin{aligned}
N_1 &= \{1248, 1268, 1348, 1378, 1568, 1578, 2358, 2378, 2458, 2678, 3468, 3568, 4578, 4678, \\
&\quad 1247, 1257, 1367, 1467, 2347, 2567, 3457, 3567, 1236, 2346, 1345, 1235, 1456, 2456\}, \\
N_2 &= \{1248, 2458, 2358, 3568, 3468, 4678, 4578, 1578, 1568, 1268, 2678, \\
&\quad 2378, 1378, 1348, 1247, 2457, 2357, 3567, 3467, 1567, 1267, 1347\} = \Sigma_{78}T, \\
N_3 &= \{1248, 1268, 1348, 1378, 1568, 1578, 2358, 2378, 2458, 2678, 3468, 3568, \\
&\quad 4578, 4678, 1234, 2347, 2456, 2467, 3456, 3457, 1235, 1256, 1357\}, \\
N_4 &= \{1248, 1268, 1348, 1378, 1568, 1578, 2358, 2378, 2458, 2678, 3468, \\
&\quad 3568, 4578, 4678, 1245, 1256, 2356, 2367, 3467, 1347, 1457\}, \\
N_5 &= \{1258, 1268, 1358, 1378, 1468, 1478, 2368, 2378, 2458, 2478, 3458, 3468, \\
&\quad 1257, 1267, 1367, 1457, 2357, 2467, 3457, 3467, 2356, 2456, 1356, 1456\}, \\
N_6 &= \{1358, 1378, 1468, 1478, 1568, 2368, 2378, 2458, 2478, 2568, 3458, 3468, \\
&\quad 1235, 1245, 1457, 1567, 2357, 2567, 3457, 1236, 1246, 1367, 2467, 3467\}, \\
N_7 &= \{1268, 1258, 1358, 1378, 1478, 1468, 2378, 2368, 2458, 2478, 3468, \\
&\quad 3458, 1356, 1367, 2357, 2356, 3467, 3457, 1256, 1467, 2457\}, \\
N_8 &= \kappa_{348}(\kappa_{238}(\kappa_{56}(\kappa_{67}(N_7)))), \quad N_9 = \kappa_{235}(\kappa_{67}(N_7)), \\
N_{10} &= \kappa_{148}(\kappa_{67}(N_7)), \quad N_{11} = \kappa_{348}(\kappa_{56}((N_{10}))), \quad N_{12} = \kappa_{457}(\kappa_{23}((N_9))), \\
N_{13} &= \kappa_{567}(\kappa_{23}((N_9))), \quad N_{14} = \kappa_{138}(\kappa_{57}((N_8))) \cong \Sigma_{78}R_2, \quad N_{15} = \kappa_{158}(\kappa_{23}(N_9)).
\end{aligned}$$

All the vertices of N_1 are singular and their links are isomorphic to the 7-vertex torus T . There are two singular vertices in N_2 and their links are isomorphic to T . The singular vertices in N_3 are 8, 3, 4, 2, 5 and their links are isomorphic to T , R_2 , R_2 , R_3 and R_3 respectively. There is only one singular vertex in N_4 whose link is isomorphic to T . All the vertices of N_5 (respectively, N_6) are singular and their links are isomorphic to R_4 (respectively, R_3). Each of $N_7, \dots, N_{15}$ has exactly two singular vertices and their links are 7-vertex $\mathbb{R}P^2$'s. Thus, each N_i is a normal 3-pseudomanifold.

It follows from the definition that $N_i \approx N_j$ for $7 \leq i, j \leq 15$. Here we prove.

Lemma 3.3. (a) The geometric carriers of N_1, N_2, N_3, N_4, N_5 and N_7 are distinct (non-homeomorphic), (b) $N_i \not\cong N_j$ for $1 \leq i < j \leq 7$, (c) $N_5 \sim N_6$.

Proof. For a normal 3-pseudomanifold X , let $n_s(X)$ denote the number of singular vertices. Clearly, if M and N are two normal 3-pseudomanifolds with homeomorphic geometric carriers then $(n_s(M), \chi(M)) = (n_s(N), \chi(N))$. Now, $(n_s(N_1), \chi(N_1)) = (8, 8)$, $(n_s(N_2), \chi(N_2)) = (2, 2)$, $(n_s(N_3), \chi(N_3)) = (5, 3)$, $(n_s(N_4), \chi(N_4)) = (1, 1)$, $(n_s(N_5), \chi(N_5)) = (8, 4)$, $(n_s(N_7), \chi(N_7)) = (2, 1)$. This proves (a).

Part (b) follows from the fact that N_i is neighbourly and has no removable edge and hence there is no proper bistellar move from N_i , for $1 \leq i \leq 6$.

Let N'_5 be obtained from N_5 by starring a new vertex 0 in the facet 1358. Let $N''_5 = \kappa_{\{0\}}(\kappa_{08}(\kappa_{156}(\kappa_{07}(\kappa_{03}(\kappa_{035}(\kappa_{68}(\kappa_{02}(\kappa_{268}(\kappa_{13}(\kappa_{135}(\kappa_{138}(\kappa_{158}(N'_5))))))))))))))$. Then N''_5 is isomorphic to N_6 via the map $(2, 3)(5, 8)$. This proves (c). $\square$

Lemma 3.4. $N_k \not\cong N_l$ for $1 \leq k < l \leq 15$.

Proof. Let n_s be as above. Clearly, if M and N are two isomorphic 3-pseudomanifolds then $(n_s(M), f_3(M)) = (n_s(N), f_3(N))$. Now, $(n_s(N_1), f_3(N_1)) = (8, 28)$, $(n_s(N_2), f_3(N_2)) = (2, 22)$, $(n_s(N_3), f_3(N_3)) = (5, 23)$, $(n_s(N_4), f_3(N_4)) = (1, 21)$, $(n_s(N_5), f_3(N_5)) = (n_s(N_6), f_3(N_6)) = (8, 24)$, and $(n_s(N_i), f_3(N_i)) = (2, 21)$ for $7 \leq i \leq 15$. Since the links of each vertex in N_5 is isomorphic to R_4 and the links of each vertex in N_6 is isomorphic to R_3 , it follows that $N_5 \not\cong N_6$. Thus, $N_i \not\cong N_j$ for $1 \leq i \leq 6$, $1 \leq j \leq 15$, $i \neq j$.

Observe that the singular vertices in N_i are 3 and 8 for $7 \leq i \leq 15$. Moreover, (i) $\text{lk}_{N_7}(3) \cong \text{lk}_{N_7}(8) \cong R_4$, (ii) $\text{lk}_{N_8}(3) \cong R_4$ and $\text{lk}_{N_8}(8) \cong R_3$, (iii) $\text{lk}_{N_9}(3) \cong R_2$ and $\text{lk}_{N_9}(8) \cong R_4$, (iv) $\text{lk}_{N_{10}}(3) \cong \text{lk}_{N_{10}}(8) \cong R_3$ and $\deg_{N_{10}}(38) = 6$, (v) $\text{lk}_{N_{11}}(3) \cong \text{lk}_{N_{11}}(8) \cong R_3$ and $\deg_{N_{11}}(38) = 5$, (vi) $\text{lk}_{N_{12}}(3) \cong R_2$, $\text{lk}_{N_{12}}(8) \cong R_3$ and $G_3(N_{12}) = (V, \{32, 21, 17, 75, 54, 46\})$, (vii) $\text{lk}_{N_{13}}(3) \cong R_2$, $\text{lk}_{N_{13}}(8) \cong R_3$ and $G_3(N_{13}) = (V, \{32, 21, 17, 75, 56, 67, 64, 42\})$, (viii) $\text{lk}_{N_{14}}(3) \cong \text{lk}_{N_{14}}(8) \cong R_2$ and $\deg_{N_{14}}(38) = 3$, (xi) $\text{lk}_{N_{15}}(3) \cong \text{lk}_{N_{15}}(8) \cong R_2$ and $\deg_{N_{15}}(38) = 6$. These imply that there is no isomorphism between N_i and N_j for $7 \leq i < j \leq 15$. This completes the proof. $\square$

Example 5: Some 8-vertex non-neighbourly normal 3-pseudomanifolds.

$$\begin{aligned} N_{16} &= \kappa_{67}(N_7), & N_{17} &= \kappa_{24}(N_8), & N_{18} &= \kappa_{238}(\kappa_{56}(\kappa_{67}(N_7))), & N_{19} &= \kappa_{57}(N_8), \\ N_{20} &= \kappa_{56}(N_{10}), & N_{21} &= \kappa_{12}(N_9), & N_{22} &= \kappa_{14}(N_{11}), & N_{23} &= \kappa_{23}(N_9), \\ N_{24} &= \kappa_{38}(N_{14}), & N_{25} &= \kappa_{56}(N_{16}), & N_{26} &= \kappa_{12}(N_{16}), & N_{27} &= \kappa_{56}(N_{17}), \\ N_{28} &= \kappa_{57}(N_{18}), & N_{29} &= \kappa_{15}(N_{18}), & N_{30} &= \kappa_{12}(N_{23}), & N_{31} &= \kappa_{24}(N_{22}), \\ N_{32} &= \kappa_{24}(N_{26}), & N_{33} &= \kappa_{57}(N_{25}), & N_{34} &= \kappa_{45}(N_{28}), & N_{35} &= \kappa_{58}(N_{29}). \end{aligned}$$

Lemma 3.5. (a) $N_i \not\cong N_j$ for $1 \leq i < j \leq 35$ and (b) $N_k \approx N_l$, for $7 \leq k, l \leq 35$.

Proof. For $0 \leq i \leq 3$, let $\mathcal{N}_i$ denote the set of 3-pseudomanifolds defined in Examples 4 and 5 with i non-edges. Then $\mathcal{N}_0 = \{N_1, \dots, N_{15}\}$, $\mathcal{N}_1 = \{N_{16}, \dots, N_{24}\}$, $\mathcal{N}_2 = \{N_{25}, \dots, N_{31}\}$ and $\mathcal{N}_3 = \{N_{32}, \dots, N_{35}\}$. The singular vertices in N_i are 3 and 8 for $7 \leq i \leq 35$.

By Lemma 3.4, the members of $\mathcal{N}_0$ are pairwise non-isomorphic.

Observe that (i) $\text{lk}_{N_{16}}(3) \cong R_4$ and $\text{lk}_{N_{16}}(8) \cong R_3$, (ii) $\text{lk}_{N_{17}}(3) \cong \text{lk}_{N_{17}}(8) \cong R_4$, (iii) $\text{lk}_{N_{18}}(3) \cong \text{lk}_{N_{18}}(8) \cong R_3$ and $G_6(N_{18}) = (V, \{73, 31, 18, 84\})$, (iv) $\text{lk}_{N_{19}}(3) \cong \text{lk}_{N_{19}}(8) \cong R_3$ and $G_6(N_{19}) = (V, \{63, 31, 18, 86\})$, (v) $\text{lk}_{N_{20}}(3) \cong \text{lk}_{N_{20}}(8) \cong R_3$ and $G_6(N_{20}) = (V, \{74, 28, 83, 31\})$, (vi) $\text{lk}_{N_{21}}(3) \cong R_2$, $\text{lk}_{N_{21}}(8) \cong R_3$ and $G_6(N_{21}) = (V, \{48, 83, 37, 36\})$,

(vii) $\text{lk}_{N_{22}}(3) \cong R_2$, $\text{lk}_{N_{22}}(8) \cong R_3$ and $G_6(N_{22}) = (V, \{28, 86, 63, 37, 38\})$, (viii) $\text{lk}_{N_{23}}(3) \cong R_1$ and $\text{lk}_{N_{23}}(8) \cong R_3$, (ix) $\text{lk}_{N_{24}}(3) \cong \text{lk}_{N_{24}}(8) \cong R_1$. These imply that there is no isomorphism between any two members of $\mathcal{N}_1$.

Observe that (i) $\text{lk}_{N_{25}}(3) \cong R_3$ and $\text{lk}_{N_{25}}(8) \cong R_4$, (ii) $\text{lk}_{N_{26}}(3) \cong \text{lk}_{N_{26}}(8) \cong R_3$ and $G_6(N_{26}) = (V, \{53, 38, 84\})$, (iii) $\text{lk}_{N_{27}}(3) \cong \text{lk}_{N_{27}}(8) \cong R_3$, $G_6(N_{27}) = (V, \{78, 81, 13, 37\})$ and $\text{NEG}(N_{27}) = \{24, 56\}$, (iv) $\text{lk}_{N_{28}}(3) \cong \text{lk}_{N_{28}}(8) \cong R_3$, $G_6(N_{28}) = (V, \{18, 84, 43, 31\})$ and $\text{NEG}(N_{28}) = \{75, 56\}$, (v) $\text{lk}_{N_{29}}(3) \cong R_3$ and $\text{lk}_{N_{29}}(8) \cong R_2$, (vi) $\text{lk}_{N_{30}}(3) \cong R_1$ and $\text{lk}_{N_{30}}(8) \cong R_3$, (vii) $\text{lk}_{N_{31}}(3) \cong \text{lk}_{N_{31}}(8) \cong R_2$. These imply that there is no isomorphism between any two members of $\mathcal{N}_2$.

Observe that (i) $\text{lk}_{N_{32}}(3) \cong \text{lk}_{N_{32}}(8) \cong R_3$, (ii) $\text{lk}_{N_{33}}(3) \cong \text{lk}_{N_{33}}(8) \cong R_4$, (iii) $\text{lk}_{N_{34}}(3) \cong \text{lk}_{N_{34}}(8) \cong R_2$, (iv) $\text{lk}_{N_{35}}(3) \cong R_2$ and $\text{lk}_{N_{35}}(8) \cong R_1$. These imply that there is no isomorphism between any two members of $\mathcal{N}_3$.

Since a member of $\mathcal{N}_i$ is non-isomorphic to a member of $\mathcal{N}_j$ for $i \neq j$, the above imply Part (a). Part (b) follows from the definition of N_k for $8 \leq k \leq 35$. $\square$

The 3-dimensional *Kummer variety* K^3 is the torus $S^1 \times S^1 \times S^1$ modulo the involution $\sigma : x \mapsto -x$. It has 8 singular points corresponding to 8 elements of order 2 in the abelian group $S^1 \times S^1 \times S^1$. In [11], Kühnel showed that N_5 triangulates K^3 . For a topological space X , $C(X)$ denotes a cone with base X . Let $H = D^2 \times S^1$ denote the solid torus. As a consequence of the above lemmas we get.

Corollary 5. *All the 8-vertex normal 3-pseudomanifolds triangulate seven distinct topological spaces, namely, $|S_{8,j}^3| = S^3$ for $1 \leq j \leq 38$, $|N_1|$, $|N_2| = S(S^1 \times S^1)$, $|N_3|$, $|N_4| = H \cup (C(\partial H))$, $|N_5| = |N_6| = K^3$ and $|N_i| = S(\mathbb{R}P^2)$ for $7 \leq i \leq 35$.*

Proof. Let K be a 8-vertex normal 3-pseudomanifold. If K is a combinatorial 3-sphere then it triangulates the 3-sphere S^3 .

If K is not a combinatorial 3-sphere then, by Lemma 3.5 (b), $|K|$ is (pl) homeomorphic to $|N_1|, \dots, |N_6|$ or $|N_7|$. Since $N_2 = \Sigma_{78}T$, $|N_2|$ is homeomorphic to the suspension $S(S^1 \times S^1)$. In N_4 , the facets not containing the vertex 8 form a solid torus whose boundary is the link of 8. This implies that $|N_4| = H \cup (C(\partial H))$. It follows from Lemma 3.3 (c) that $|N_6|$ is (pl) homeomorphic to $|N_5| = K^3$. Since N_{24} is isomorphic to the suspension $S_2^0 * R_1$, $|N_{24}| = S(\mathbb{R}P^2)$. Therefore, by Lemma 3.5 (b), $|N_i|$ is (pl) homeomorphic to $|N_{24}| = S(\mathbb{R}P^2)$ for $7 \leq i \leq 35$. The result now follows from Lemma 3.3 (a). $\square$

A 3-dimensional *pseudo-complex* K is an ordered pair (Δ, Φ) , where Δ is a finite collection of disjoint tetrahedra and Φ is a family of affine isomorphisms between pairs of 2-faces of the tetrahedra in Δ . Let $|K|$ denote the quotient space obtained from the disjoint union $\sqcup_{\sigma \in \Delta} \sigma$ by setting $x = \varphi(x)$ for $\varphi \in \Phi$. The quotient of a tetrahedron $\sigma \in \Delta$ in $|K|$ is called a *3-simplex* in $|K|$ and is denoted by $|\sigma|$. Similarly, the quotient of 2-faces, edges and vertices of tetrahedra are called *2-simplices*, *edges* and *vertices* in $|K|$ respectively. If $|K|$ is homeomorphic to a topological space X then K is called a *pseudo-triangulation* of X . A 3-dimensional pseudo-complex $K = (\Delta, \Phi)$ is said to be *regular* if (i) each 3-simplex in $|K|$ has four distinct vertices and (ii) for $2 \leq i \leq 3$, no two distinct i -simplices in $|K|$ have the same set of vertices. So, for $2 \leq i \leq 3$, an i -simplex α in $|K|$ is uniquely determined by its vertices and denoted by $u_1 \cdots u_{i+1}$, where $u_1, \dots, u_{i+1}$ are vertices of α . (But, the edges in $|K|$ may not form a simple graph.) So, we can identify a regular pseudo-complex $K = (\Delta, \Phi)$ with $\mathcal{K} := \{|\sigma| : \sigma \in \Delta\}$. Simplices and edges in $|K|$ are said to be simplices and edges of $\mathcal{K}$. Clearly, a pure 3-dimensional simplicial complex is a regular pseudo-complex.

Let $\mathcal{M}$ be a regular pseudo-triangulation of X and $abcd, abce$ be two 3-simplices in $\mathcal{M}$. If ade, bde, cde are not 2-simplices in $\mathcal{M}$, then $\mathcal{N} := (\mathcal{M} \setminus \{abcd, abce\}) \cup \{abde, acde, bcde\}$ is also a regular pseudo-triangulation of X . We say that $\mathcal{N}$ is obtained from $\mathcal{M}$ by the *generalized bistellar 1-move* κ_{abc} . If there is no edge between d and e in $\mathcal{M}$ then κ_F is called a *bistellar 1-move*. If there exist 3-simplices of the form $xyuv, xzuv, yzuv$ in a regular pseudo-triangulation $\mathcal{P}$ of Y and xyz is not a 2-simplex, then $\mathcal{Q} := (\mathcal{P} \setminus \{xyuv, xzuv, yzuv\}) \cup \{xyzu, xyzv\}$ is also a regular pseudo-triangulation of Y . We say that $\mathcal{Q}$ is obtained from $\mathcal{P}$ by the *generalized bistellar 2-move* κ_E , where E is the common edge in $xyuv, xzuv$ and $yzuv$. If E is the only edge between u and v in $\mathcal{P}$ then κ_E is called a *bistellar 2-move*.

Let M be a pseudo-triangulation of a closed 3-manifold and N a 3-pseudomanifold. A simplicial map $f: M \rightarrow N$ is said to be a *k-fold branched covering* (with discrete branch locus) if there exists $U \subseteq V(N)$ such that $|f|_{|M| \setminus f^{-1}(U)}: |M| \setminus f^{-1}(U) \rightarrow |N| \setminus U$ is a *k-fold covering*. The smallest such U (so that $|f|_{|M| \setminus f^{-1}(U)}: |M| \setminus f^{-1}(U) \rightarrow |N| \setminus U$ is a covering) is called the *branch locus*. It is known that N_1 can be regarded as a branched quotient of a regular hyperbolic tessellation (cf. [13]). In [11], Kühnel has shown that N_5 is a 2-fold branched quotient of a pseudo-triangulation of the 3-dimensional torus. Here we prove:

Theorem 6. (a) N_{24} is a 2-fold branched quotient of a 14-vertex combinatorial 3-sphere.
(b) For $7 \leq i \leq 35$, N_i is a 2-fold branched quotient of a 14-vertex regular pseudo-triangulation of the 3-sphere.

Lemma 3.6. Let M be a regular pseudo-triangulation of a 3-manifold and N be a normal 3-pseudomanifold. Let $f: M \rightarrow N$ be a *k-fold branched covering* with at most two vertices in the branch locus. If $\kappa_e: N \mapsto \tilde{N}$ is a bistellar 2-move, then there exist *k* generalized bistellar 2-moves $\kappa_{e_1}, \dots, \kappa_{e_k}$ such that $\kappa_{e_k}(\dots(\kappa_{e_1}(M)))$ is a *k-fold branched cover* of $\tilde{N}$.

Proof. Let $\text{lk}_N(e) = S_3^1(\{x, y, z\})$. Let $f^{-1}(e)$ consists of the edges $e_1, \dots, e_k$. Let the end points of e_i be u_i, v_i , the 3-simplices containing e_i be $u_i v_i x_i y_i, u_i v_i x_i z_i, u_i v_i y_i z_i$ and $f(x_i) = x, f(y_i) = y, f(z_i) = z$ for $1 \leq i \leq k$. Since xyz is not a simplex in N , it follows that $x_i y_i z_i$ is not a 2-simplex in M . Let M_i be the pseudo-complex consists of $u_i v_i x_i y_i, u_i v_i x_i z_i$ and $u_i v_i y_i z_i$. Since the number of vertices in the branched locus is at most 2, it follows that the number of vertices common in M_i and M_j is at most 2 for $i \neq j$. In particular, $\#(\{x_i, y_i, z_i\} \cap \{x_j, y_j, z_j\}) \leq 2$. Therefore, $x_j y_j z_j$ is not a 2-simplex in $\kappa_{e_i}(M)$. So, we can perform generalized bistellar 2-move κ_{e_j} on $\kappa_{e_i}(M) = (M \setminus M_i) \cup \{x_i y_i z_i u_i, x_i y_i z_i v_i\}$ for $i \neq j$. Clearly, $\tilde{M} := \kappa_{e_k}(\dots \kappa_{e_1}(M))$ is a *k-fold branched cover* of $\tilde{N}$ (via the map $\tilde{f}$, where $\tilde{f}(w) = f(w)$ for $w \in V(\tilde{M}) = V(M)$ and $\tilde{f}(x_i y_i z_i u_i) = xyzu$ and $\tilde{f}(x_i y_i z_i v_i) = xyzv$). $\square$

Lemma 3.7. Let M be a regular pseudo-triangulation of a 3-manifold and N be a normal 3-pseudomanifold. Let $f: M \rightarrow N$ be a *k-fold branched covering* with at most two vertices in the branch locus. If $\kappa_F: N \mapsto \tilde{N}$ is a bistellar 1-move, then there exist *k* generalized bistellar 1-moves $\kappa_{F_1}, \dots, \kappa_{F_k}$ such that $\kappa_{F_k}(\dots(\kappa_{F_1}(M)))$ is a *k-fold branched cover* of $\tilde{N}$.

Proof. Let $F = xyz$ and $\text{lk}_N(F) = \{u, v\}$. Let $f^{-1}(F)$ consist of the 2-simplices $F_1, \dots, F_k$. Let $F_i = x_i y_i z_i$ and the 3-simplices containing F_i be $x_i y_i z_i u_i$ and $x_i y_i z_i v_i$ and $f(x_i, y_i, z_i, u_i, v_i) = (x, y, z, u, v)$ for $1 \leq i \leq k$. Since f is simplicial, it follows that $x_i u_i v_i, y_i u_i v_i$ and $z_i u_i v_i$ are not 2-simplices in M . Let M_i be pseudo-complex $\{x_i y_i z_i u_i, x_i y_i z_i v_i\}$. Since the number of vertices in the branched locus is at most 2, it follows that $x_j u_j v_j, y_j u_j v_j$ and $z_j u_j v_j$ are not 2-simplices in $\kappa_{F_i}(M)$ for $i \neq j$. Then (by the similar arguments as in the proof of Lemma 3.6) $\kappa_{F_k}(\dots \kappa_{F_1}(M))$ is a *k-fold branched cover* of $\tilde{N}$. $\square$

Proof of Theorem 6. If $\mathcal{I}$ denotes the boundary of the icosahedron then there exists a simplicial 2-fold covering $f: \mathcal{I} \rightarrow R_1$. Consider the simplicial map $\tilde{f}: S_2^0(\{a, b\}) * \mathcal{I} \rightarrow S_2^0(\{c, d\}) * R_1$ given by $\tilde{f}(a) = c$, $\tilde{f}(b) = d$ and $\tilde{f}(u) = f(u)$ for $u \in V(\mathcal{I})$. Then $\tilde{f}$ is a 2-fold branched covering with branch locus $\{c, d\}$. Since N_{24} is isomorphic to the suspension $S_2^0 * R_1$, it follows that N_{24} is a 2-fold branched quotient of the 14-vertex combinatorial 3-sphere $S_2^0(\{a, b\}) * \mathcal{I}$ (with branch locus $\{3, 8\}$). This proves (a).

The result now follows from Lemmas 3.5 (a), 3.6 and 3.7. (In fact, to obtain a 2-fold branched cover $\tilde{N}_{14}$ of N_{14} from $R_1 * S_2^0$ one needs one bistellar 1-move and then one generalized bistellar 1-move. And all other moves required in the proof are bistellar moves on regular pseudo-triangulations of S^3 .) $\square$

Remark 1. The combinatorial 3-sphere $R_1 * S_2^0$ is a 2-fold branched cover of N_{24} and N_{14} can be obtained from N_{24} by a bistellar 1-move. Now, if $f: M \rightarrow N_{14}$ is a 2-fold branched covering and M is a combinatorial 3-manifold then (since $\text{lk}_{N_{14}}(8)$ is a 7-vertex triangulated $\mathbb{R}P^2$) the link of any vertex in $f^{-1}(8)$ is a 14-vertex triangulated S^2 and hence $f_0(M) > 14$. (Similarly, for $i \neq 24$, if N_i is a branched quotient of a combinatorial 3-manifold M then $f_0(M) > 14$.) So, there does not exist a combinatorial 3-sphere M which is a branched cover of N_{14} and which can be obtained from $R_1 * S_2^0$ by proper bistellar moves.

In [2], Altshuler observed that N_1 is orientable and $|N_1|$ is simply-connected. In [14], Lutz showed that $(H_1(N_1), H_2(N_1), H_3(N_1)) = (0, \mathbb{Z}^8, \mathbb{Z})$. The normal 3-pseudomanifold N_3 is the only among all the 35 which has singular vertices of different types, namely one singular vertex whose link is a triangulated torus and four singular vertices whose links are triangulated real projective planes. Using polymake ([9]), we find that $(H_1(N_3), H_2(N_3), H_3(N_3)) = (0, \mathbb{Z}^2 \oplus \mathbb{Z}_2, 0)$. We summarized all the findings about $N_1, \dots, N_{35}$ in the next table.

Table 1 : 8-vertex normal 3-pseudomanifolds which are not combinatorial 3-manifolds.

X	f -vector (f_1, f_2, f_3)	$\chi(X)$	$n_s(X)$	links of singular vertices	Geometric carriers, Homology (H_1, H_2, H_3)
N_1	(28, 56, 28)	8	8	all are T	$ N_1 $ is simply connected, (H_1, H_2, H_3) = $(0, \mathbb{Z}^8, \mathbb{Z})$
N_2	(28, 44, 22)	2	2	both are T	$ N_2 = S(S^1 \times S^1)$
N_3	(28, 46, 23)	3	5	$T, R_2, R_2,$ R_3, R_3	$(H_1, H_2, H_3) =$ $(0, \mathbb{Z}^2 \oplus \mathbb{Z}_2, 0)$
N_4	(28, 42, 21)	1	1	T	$ N_4 = H \cup (C(\partial H))$
N_5	(28, 48, 24)	4	8	all are R_4	$ N_5 = K^3$
N_6	„	„	„	all are R_3	$ N_6 = K^3$
N_7	(28, 42, 21)	1	2	both are R_4	$ N_7 = S(\mathbb{R}P^2)$
$N_i, 8 \leq i \leq 15$	„	„	„	both are in $\{R_1, \dots, R_4\}$	$ N_i = S(\mathbb{R}P^2)$
$N_i, 16 \leq i \leq 24$	(27, 40, 20)	„	„	„	„
$N_i, 25 \leq i \leq 31$	(26, 38, 19)	„	„	„	„
$N_i, 32 \leq i \leq 35$	(25, 36, 18)	„	„	„	„

[Here K^3 is the 3-dimensional Kummer variety, $H = D^2 \times S^1$ is the solid torus, $S(Y)$ is the topological suspension of Y and $n_s(X)$ is the number of singular vertices in X .]

Example 6: For $d \geq 2$, let

$$K_{2d+3}^d = \{v_i \cdots v_{j-1} v_{j+1} \cdots v_{i+d+1} : i+1 \leq j \leq i+d, 1 \leq i \leq 2d+3\}$$

(additions in the suffixes are modulo $2d+3$). It was shown in [12] the following: (i) K_{2d+3}^d is a triangulated d -manifold for all $d \geq 2$, (ii) K_{2d+3}^d triangulates $S^{d-1} \times S^1$ for d even, and triangulates the twisted product $S^{d-1} \times S^1$ (the twisted S^{d-1} -bundle over S^1) for d odd. For $d \geq 3$, K_{2d+3}^d is the unique non-simply connected $(2d+3)$ -vertex triangulated d -manifold (cf. [4]). The combinatorial 3-manifolds K_9^3 was first constructed by Walkup in [15].

From K_9^3 we construct the following 10-vertex combinatorial 3-manifold.

$$\begin{aligned} A_{10}^3 := & (K_9^3 \setminus \{v_1 v_2 v_3 v_5, v_2 v_3 v_5 v_6, v_3 v_5 v_6 v_7, v_3 v_4 v_6 v_7, v_4 v_6 v_7 v_8\}) \cup \\ & \{v_0 v_1 v_2 v_3, v_0 v_1 v_2 v_5, v_0 v_1 v_3 v_5, v_0 v_2 v_3 v_6, v_0 v_2 v_5 v_6, v_0 v_3 v_5 v_7, \\ & v_0 v_5 v_6 v_7, v_0 v_3 v_4 v_6, v_0 v_3 v_4 v_7, v_0 v_4 v_6 v_8, v_0 v_4 v_7 v_8, v_0 v_6 v_7 v_8\}. \end{aligned}$$

[Geometrically, first we remove a pl 3-ball consisting of five 3-simplices from $|K_9^3|$. This gives a pl 3-manifold with boundary and the boundary is a 2-sphere. Then we add a cone with base this boundary and vertex v_0 . So, the new polyhedron $|A_{10}^3|$ is pl homeomorphic to $|K_9^3|$. This implies that the simplicial complex A_{10}^3 is a combinatorial 3-manifold.]

The only non-edge in A_{10}^3 is $v_0 v_9$ and there is no common 2-face in the links of v_0 and v_9 in A_{10}^3 . So, A_{10}^3 does not allow any bistellar 1-move. So, A_{10}^3 is a 10-vertex non-neighbourly combinatorial 3-manifold which does not admit any bistellar 1-move.

Similarly, from K_{11}^4 we construct the following 12-vertex triangulated 4-manifold.

$$\begin{aligned} A_{12}^4 := & (K_{11}^4 \setminus \{v_1 v_2 v_3 v_4 v_6, v_2 v_3 v_4 v_6 v_7, v_3 v_4 v_6 v_7 v_8, v_4 v_6 v_7 v_8 v_9, v_4 v_5 v_7 v_8 v_9, v_5 v_7 v_8 v_9 v_{10}\}) \\ & \cup \{v_0 v_1 v_2 v_3 v_4, v_0 v_1 v_2 v_3 v_6, v_0 v_1 v_2 v_4 v_6, v_0 v_1 v_3 v_4 v_6, v_0 v_2 v_3 v_4 v_7, v_0 v_2 v_3 v_6 v_7, \\ & v_0 v_2 v_4 v_6 v_7, v_0 v_3 v_4 v_6 v_8, v_0 v_3 v_4 v_7 v_8, v_0 v_3 v_6 v_7 v_8, v_0 v_4 v_6 v_7 v_9, v_0 v_4 v_6 v_8 v_9, \\ & v_0 v_4 v_7 v_8 v_9, v_0 v_4 v_5 v_7 v_9, v_0 v_4 v_5 v_8 v_9, v_0 v_4 v_7 v_8 v_9, v_0 v_5 v_7 v_8 v_{10}, v_0 v_5 v_7 v_9 v_{10}, \\ & v_0 v_5 v_8 v_9 v_{10}\}. \end{aligned}$$

The only non-edge in A_{12}^4 is $v_0 v_{11}$ and there is no common 2-face in the links of v_0 and v_{11} in A_{12}^4 . So, A_{12}^4 does not allow any bistellar 1-move. So, A_{12}^4 is a 12-vertex non-neighbourly triangulated 4-manifold which does not admit any bistellar 1-move.

By the same way, one can construct a $(2d+4)$ -vertex non-neighbourly triangulated d -manifold A_{2d+4}^d (from K_{2d+3}^d) which does not admit any bistellar 1-move for all $d \geq 3$.

Example 7: Let N_3 be as in Example 4. Let M be obtained from N_3 by starring two vertices u and v in the facets 1248 and 3568 respectively, i.e., $M = \kappa_{1248}(\kappa_{3568}(N_3))$. Then M is a 10-vertex normal 3-pseudomanifold. Let B_9^3 be obtained from M by identifying the vertices u and v . Let the new vertex be 9. Then

$$B_9^3 := (N_3 \setminus \{1248, 3568\}) \cup \{1249, 1289, 1489, 2489, 3569, 3589, 3689, 5689\}.$$

The degree 3 edges in B_9^3 are 16, 17 and 67. But, none of these edges is removable. So, no bistellar 2-moves are possible from B_9^3 . The only non-edge in B_9^3 is 79. Since there is no common 2-face in the links of 7 and 9, no bistellar 1-move is possible. So, B_9^3 is a 9-vertex non-neighbourly 3-pseudomanifold which does not admit any proper bistellar move.

4 Proofs

For $n \geq 4$, by an S_n^2 we mean a combinatorial 2-sphere on n vertices. If $\kappa_\beta: M \mapsto N$ is a bistellar 1-move then $\deg_N(v) \geq \deg_M(v)$ for $v \in V(M)$. Here we prove the following.

Lemma 4.1. *Let M be an n -vertex 3-pseudomanifold and u be a vertex of degree 4. If $n \geq 6$ then there exists a bistellar 1-move $\kappa_\beta: M \mapsto N$ such that $\deg_N(u) = 5$.*

Proof. Let $\text{lk}_M(u) = S_4^2(\{a, b, c, d\})$ and $\beta = abc$. Let $\text{lk}_M(\beta) = \{u, x\}$. If $x = d$ then the induced complex $K = M[\{u, a, b, c, d\}]$ is a 3-pseudomanifold. Since $n \geq 6$, K is a proper subcomplex of M . This is not possible. So, $x \neq d$ and hence ux is a non-edge in M . Then κ_β is a bistellar 1-move. Since ux is an edge in $\kappa_\beta(M)$, κ_β is a required bistellar 1-move. $\square$

Lemma 4.2. *Let M be an n -vertex 3-pseudomanifold and u be a vertex of degree 5. If $n \geq 7$ then there exists a bistellar 1-move $\kappa_\beta: M \mapsto N$ such that $\deg_N(u) = 6$.*

Proof. Since $\deg_M(u) = 5$, the link of u in M is of the form $S_2^0(\{a, b\}) * S_3^1(\{x, y, z\})$ for some vertices a, b, x, y, z of M . If both $xyza$ and $xuzb$ are facets then the induced subcomplex $M[\{x, y, z, u, a, b\}]$ is a 3-pseudomanifold. This is not possible since $n \geq 7$. So, without loss of generality, assume that $xyza$ is not a facet. Again, if $xyab, xzab$ and $yzab$ all are facets then the induced subcomplex $M[\{u, x, y, z, a, b\}]$ is a 3-pseudomanifold, which is not possible. So, assume that $xyab$ is not a facet.

Consider the face $\beta = xya$. Suppose $\text{lk}_M(\beta) = \{u, w\}$. From the above, $w \notin \{z, b\}$. So, uw is a non-edge and hence κ_β is a required bistellar 1-move. $\square$

Lemma 4.3. *Let M be a non-neighbourly 8-vertex 3-pseudomanifold and u be a vertex of degree 6. If the degree of each vertex is at least 6, then there exists a bistellar 1-move $\kappa_\tau: M \mapsto N$ such that $\deg_N(u) = 7$.*

Proof. Let u be a vertex with $\deg_M(u) = 6$ and uv be a non-edge. Let $L = \text{lk}_M(u)$.

Claim. There exists a 2-face τ such that $\tau \cup \{u\}$ and $\tau \cup \{v\}$ are facets.

First consider the case when there exists a vertex w such that $\deg_L(w) = 5$. Let $\text{lk}_L(w) (= \text{lk}_M(uw)) = C_5(1, 2, 3, 4, 5)$.

Let $K = \text{lk}_M(w)$. Since $\deg(v) = 6$, vw is an edge. Thus K contains 7 vertices. If one of $12v, \dots, 45v, 51v$ is a 2-face, say $12v$, then $12wv$ and $12wu$ are facets. In this case, $\tau = 12w$ serves the purpose. So, assume that $12v, \dots, 45v, 51v$ are non-faces in K . Then there are at least three 2-faces (not containing u) containing the edges $12, \dots, 45, 51$ in K . Also, there are at least three 2-faces containing v in K . So, the number of 2-faces in K is at least 11. This implies that $\deg_K(v) = 3$ or 4 and K is a 7-vertex $\mathbb{R}P^2$ or P_4 . Since $\deg_K(u) = 5$, it follows that K is isomorphic to R_2, R_3 or P_4 (defined in Section 2). In each case, (since $\deg_K(u) = 5$, $\deg_K(v) = 3$ or 4 and uv is a non-edge) there exists an edge α in K such that $\alpha \cup \{u\}$ and $\alpha \cup \{v\}$ are 2-faces in K and hence $\tau = \alpha \cup \{w\}$ serves the purpose.

Now, assume that L has no vertex of degree 5. Then L must be of the form $S_2^0(\{a_1, a_2\}) * S_2^0(\{b_1, b_2\}) * S_2^0(\{c_1, c_2\})$. If possible let $a_i b_j c_k v$ is not a facet for $1 \leq i, j, k \leq 2$. Consider the 2-face $a_1 b_1 c_1$. There exists a vertex $x \neq u$ such that $a_1 b_1 c_1 x$ is a facet. Assume, without loss of generality, that $a_1 b_1 c_1 a_2$ is a facet. Since $\deg(c_1) > 5$ (respectively, $\deg(b_1) > 5$), $a_1 a_2 b_2 c_1$ (respectively, $a_1 a_2 b_1 c_2$) is not a facet. So, the facet (other than $a_1 b_2 c_1 u$) containing $a_1 b_2 c_1$ must be $a_1 b_2 c_1 c_2$. Similarly, the facet (other than $a_1 b_1 c_2 u$) containing $a_1 b_1 c_2$ must be $a_1 b_1 b_2 c_2$. Then $a_1 b_2 c_1 c_2$, $a_1 b_1 b_2 c_2$ and $a_1 b_2 c_2 u$ are three facets containing $a_1 b_2 c_2$, a contradiction. This proves the claim.

By the claim, there exists a 2-simplex τ such that $\text{lk}_M(\tau) = \{u, v\}$. Since uv is a non-edge of M , $\kappa_\tau : M \mapsto \kappa_\tau(M) = N$ is a bistellar 1-move. Since uv is an edge in N , it follows that $\deg_N(u) = 7$. $\square$

Proof of Theorem 1. Let M be an 8-vertex 3-pseudomanifold. Then, by Lemma 4.1, there exist bistellar 1-moves $\kappa_{A_1}, \dots, \kappa_{A_k}$, for some $k \geq 0$, such that the degree of each vertex in $\kappa_{A_k}(\dots(\kappa_{A_1}(M)))$ is at least 5. Therefore, by Lemma 4.2, there exist bistellar 1-moves $\kappa_{A_{k+1}}, \dots, \kappa_{A_l}$, for some $l \geq k$, such that the degree of each vertex in $\kappa_{A_l}(\dots\kappa_{A_k}(\dots(\kappa_{A_1}(M))))$ is at least 6. Then, by Lemma 4.3, there exist bistellar 1-moves $\kappa_{A_{l+1}}, \dots, \kappa_{A_m}$, for some $m \geq l$, such that the degree of each vertex in $\kappa_{A_m}(\dots\kappa_{A_l}(\dots\kappa_{A_k}(\dots(\kappa_{A_1}(M))))$ is 7. This proves the theorem. $\square$

Lemma 4.4. *Let K be an 8-vertex combinatorial 3-manifold. If K is neighbourly then K is isomorphic to $S_{8,35}^3$, $S_{8,36}^3$, $S_{8,37}^3$ or $S_{8,38}^3$.*

Proof. Since K is a neighbourly combinatorial 3-manifold, by Proposition 2.1, the link of any vertex is isomorphic to $S_5, \dots, S_8$ or S_9 .

Claim: The links of all the vertices can not be isomorphic to S_9 ($= S_2^0 * C_5$).

Otherwise, let $\text{lk}(8) = S_2^0(6, 7) * C_5(1, 2, \dots, 5)$. Consider the vertex 2. Since the degree of 2 is 7, 1267 or 2367 is not a facet. Assume without loss of generality that 1267 is not a facet. Again, if 1236 is a facet then $\deg_{\text{lk}(2)}(6) = 3$ and hence $\text{lk}(2) \not\cong S_9$. So, 1236 is not a facet. Similarly, 1256 is not a facet. Then the facet other than 1268 containing 126 must be 1246. Similarly, 1247 is a facet. This implies that $\text{lk}(2) = S_2^0(6, 7) * C_5(1, 4, 5, 3, 8)$. Thus $\deg(26) = 5$. Similarly, $\deg(16) = \deg(36) = \deg(46) = \deg(56) = 5$. Then, the 7-vertex 2-sphere $\text{lk}(6)$ contains five vertices of degree 5. This is not possible. This proves the claim.

Case 1. Consider the case when K has a vertex, (say 8) whose link is isomorphic to S_8 . Assume, without loss of generality, that the facets containing the vertex 8 are 1238, 1268, 1348, 1458, 1568, 2348, 2478, 2678, 4578 and 5678. Since $\deg(3) = 7$, $1234 \notin K$. Hence the facet other than 1238 containing the face 123 is one of 1235, 1236 or 1237.

If $1236 \in K$ then, clearly, $\deg(17) = 3$ or 4. If $\deg(17) = 4$, then on completing $\text{lk}(1)$, we see that $1457, 1567 \in K$, thereby showing that $\deg(5) = 5$, an impossibility. Hence, $\deg(17) = 3$ and therefore $1457 \in K$. There are two possibilities for the completion of $\text{lk}(1)$. If $1347, 1356, 1357 \in K$, from the links of 4 and 3, we see that $2346, 2467, 3467, 3567 \in K$. Here, $\deg(5) = 6$. If $1346, 1467, 1567 \in K$, then $\deg(5) = 5$. Thus, $1236 \notin K$.

Subcase 1.1. $1235 \in K$. Since $\deg(1) = 7$, either 1345 or 1256 is a facet. In the first case, $1257, 1267, 1567 \in K$. Here, $\deg(6) = 5$, a contradiction. So, $1256 \in M$ and hence $1347, 1357, 1457 \in K$. From the links of the vertices 1, 4, 7 and 5 we see that $1256, 2346, 2467, 3467, 3567, 2356 \in K$. Here, $K \cong S_{8,38}^3$ by the map $(1, 5, 8, 6)(2, 7)(3, 4)$.

Subcase 1.2. $1237 \in K$. By the same argument as in Subcase 1.1 (replace the vertex 1 by vertex 2) we get $1267, 2345, 2357, 2457 \in K$. From $\text{lk}(1)$ and $\text{lk}(7)$, $1346, 1456, 3456, 1367, 3567 \in K$. Here, $K \cong S_{8,38}^3$ by the map $(1, 7, 8, 6)(2, 5)(3, 4)$.

Case 2. K has no vertex whose link is isomorphic to S_8 but has a vertex whose link is isomorphic to S_6 . Using the same method as in Subcase 1.1, we find that $K \cong S_{8,37}^3$.

Case 3. K has no vertex whose link is isomorphic to S_8 or S_6 but has a vertex whose link is isomorphic to S_7 . Using the same method as in Subcase 1.1, we find that $K \cong S_{8,36}^3$.

Case 4. K has no vertex whose link is isomorphic to S_6, S_7 or S_8 but has a vertex (say 8) whose link is isomorphic to S_5 . The facets through 8 can be assumed to be 1238, 1278, 1348, 1458, 1568, 1678, 2348, 2458, 2568 and 2678. Clearly, $1234, 1267 \notin K$. If $\deg(15) = 6$, then

from $\text{lk}(1)$ and $\text{lk}(5)$, we see that $1235, 1345, 2345 \in K$, thereby showing that $\deg(3) = 5$. Hence $1237 \in K$. Now, we can assume without loss of generality that the facets required to complete $\text{lk}(1)$ are $1347, 1457$ and 1567 . Now, consider $\text{lk}(2)$. If $\deg(27) = 6$, then after completing the links of 2 and 7 we observe that $\deg(4) = 6$. Hence $\deg(23) = 6$. The links of 2, 7 and 6 show that $2345, 2356, 2367, 3467, 4567$ and $3456 \in K$. Here, $K \cong S_{8,35}^3$ by the map $(2, 3, 4, 5, 6, 7, 8)$. This completes the proof. $\square$

Lemma 4.5. *Let K be an 8-vertex neighbourly normal 3-pseudomanifold. If K has one vertex whose link is the 7-vertex torus T then K is isomorphic to N_1, N_2, N_3 or N_4 .*

Proof. Let us assume that $V(K) = \{1, \dots, 8\}$ and the link of the vertex 8 is the 7-vertex torus T . So, the facets containing 8 are $1248, 1268, 1348, 1378, 1568, 1578, 2358, 2378, 2458, 2678, 3468, 3568, 4578$ and 4678 . We have the following cases.

Case 1. There is a vertex (other than the vertex 8), say 7, whose link is isomorphic to T . Then $\text{lk}(7)$ has no vertex of degree 3 and hence $2367, 1457, 1237, 1357 \notin K$. This implies that the facet (other than 1378) containing 137 is 1367 or 1347 . In the first case, $\text{lk}(17) = C_6(5, 8, 3, 6, 4, 2)$. Thus, $1367, 1467, 1247, 1257 \in K$. Then, from the links of 67 and 37, we get $2567, 3567, 2347, 3457 \in K$. Now, from $\text{lk}(34)$, $1346 \notin K$. Then, from the links of 36, 34, 23, 14 and 26, we get $1236, 2346, 1345, 1235, 1456, 2456 \in K$. Here, $K = N_1$.

In the second case, $\text{lk}(37) = C_6(2, 8, 1, 4, 6, 5)$. Thus, $1347, 3467, 3567, 2357 \in K$. Now, from the links of 47 and 67, we get $1247, 2457, 1567, 1267 \in K$. Here, $K = N_2$.

Case 2. There is a vertex whose link is a 7-vertex $\mathbb{R}P^2$.

Claim : There exists a vertex in K whose link is isomorphic to R_2 .

If there is vertex whose link is isomorphic to R_2 then we are done. Otherwise, since $\text{Aut}(\text{lk}(8))$ acts transitively on $\{1, \dots, 7\}$, assume that $\text{lk}(4) \cong R_3$ (respectively, R_4). Since $(1, 2, 5, 7, 6, 3) \in \text{Aut}(\text{lk}(8))$, we may assume that the degree 4 vertex (respectively vertices) in $\text{lk}(4)$ is 1 (respectively are 1, 5, 6). Then, from $\text{lk}(4)$, $1247, 1347, 2467 \in K$. This implies that $\text{lk}(7)$ is a non-sphere and $\deg(67) = 3$. Hence $\text{lk}(7) \cong R_2$. This proves the claim.

By the claim, we can assume that $\text{lk}(4) \cong R_2$. Again, we may assume that the vertex 1 is of degree 3 in $\text{lk}(4)$. Then, from $\text{lk}(4)$, $1234, 2347, 2456, 2467, 3456, 3457 \in K$. Considering the links of the edges 36, 26, 27, 25 and 13, we get $1256, 1235, 1357 \in K$. Here, $K = N_3$.

Case 3. Only singular vertex in K is 8. So, the link of each vertex (other than vertex 8) is an S_7^2 (a 7-vertex 2-sphere). Since 8 is a degree 6 vertex in $\text{lk}(u)$, it follows that $\text{lk}(u)$ is isomorphic to one of S_5, S_6 or S_7 (defined in Example 1) for any vertex $u \neq 8$. If $\text{lk}(1) \cong S_5$, then (since $(3, 4, 2, 6, 5, 7) \in \text{Aut}(\text{lk}(8))$), we may assume that the other degree 6 vertex in $\text{lk}(1)$ is 3. Then, from the links of 1 and 3, $1348, 1234, 1346$ are facets containing 134, a contradiction. If $\text{lk}(1) \cong S_6$ then (since $\text{lk}(18) = C_6(3, 4, 2, 6, 5, 7)$), we may assume that the degree 5 vertices in $\text{lk}(1)$ are 2, 3, and 5. Then $\text{lk}(3)$ can not be an S_7^2 , a contradiction. So, $\text{lk}(1) \cong S_7$. Since $\text{Aut}(\text{lk}(8))$ acts transitively on $\{1, \dots, 7\}$, it follows that the link of each vertex is isomorphic to S_7 .

Since $\text{lk}(18) = C_6(3, 4, 2, 6, 5, 7)$ and $(3, 4, 2, 6, 5, 7) \in \text{Aut}(\text{lk}(8))$, we may assume that the degree 5 vertices in $\text{lk}(1)$ are 4 and 5. Since $\text{lk}(4) \cong S_7$, it follows that $1456 \notin K$. Then, from $\text{lk}(1)$, $1245, 1256, 1347, 1457 \in K$. Now, from the links of 4 and 5, we get $3467, 2356 \in K$. Then, from $\text{lk}(2)$, $2367 \in K$. Here $K = N_4$. This completes the proof. $\square$

Lemma 4.6. *Let K be an 8-vertex neighbourly normal 3-pseudomanifold. If K is not a combinatorial 3-manifold and has no vertex whose link is isomorphic to the 7-vertex torus T then K is isomorphic to $N_5, \dots, N_{14}$ or N_{15} .*

Proof. Let n_s be the number of singular vertices in K . Since K is neighbourly, by Proposition 2.2, the link of any vertex is either a 7-vertex $\mathbb{R}P^2$ or a 7-vertex S^2 . So, the number of facets through a singular (respectively non-singular) vertex is 12 (respectively 10). Let f_3 be the number of facets of K . Consider the set $S = \{(v, \sigma) : \sigma \text{ is a facet of } K \text{ and } v \in \sigma \text{ is a vertex}\}$. Then $f_3 \times 4 = \#(S) = n_s \times 12 + (8 - n_s) \times 10 = 80 + 2n_s$. This implies n_s is even. Since K is not a combinatorial 3-manifold, it follows that $n_s \neq 0$ and hence $n_s \geq 2$. So, K has at least two vertices whose links are isomorphic to R_2 , R_3 or R_4 .

Case 1. There exist (at least) two vertices whose links are isomorphic to R_4 . Assume that $\text{lk}_M(8) = R_4$. Then 1258, 1268, 1358, 1378, 1468, 1478, 2368, 2378, 2458, 2478, 3458, 3468 $\in K$. Since $(1, 3, 4)(5, 6, 7)$, $(1, 2)(3, 4) \in \text{Aut}(\text{lk}(8))$, we may assume that $\text{lk}(3)$ or $\text{lk}(7) \cong R_4$.

Subcase 1.1. $\text{lk}(7) \cong R_4$. Since $\text{lk}_{\text{lk}(7)}(8) = C_4(1, 3, 2, 4)$, it follows that 1, 2, 3, 4 are degree 5 vertices in $\text{lk}(7)$. Since $(3, 4)(5, 6) \in \text{Aut}(\text{lk}(8))$, assume without loss that 136, 145 $\in \text{lk}(7)$. Then, from $\text{lk}(7)$, we get 1257, 1267, 1367, 1457, 2357, 2467, 3457, 3467 $\in K$. This shows that $\text{lk}(2)$ is an $\mathbb{R}P_7^2$. Since 3457, 3458 $\in K$, it follows that 2345 $\notin K$. Then, from $\text{lk}(2)$, 2356, 2456 $\in K$. Then, from the links of 3 and 4, 1356, 1456 $\in K$. Here $K = N_5$.

Subcase 1.2. $\text{lk}(7) \not\cong R_4$. So, $\text{lk}(3) \cong R_4$. Since $\text{lk}_{\text{lk}(3)}(8) = C_6(1, 7, 2, 6, 4, 5)$, the degree 4 vertices in $\text{lk}(3)$ are either 5, 6, 7 or 1, 2, 4. In the first case, on completion of $\text{lk}(3)$, we observe that 56, 67, 57 remain non-edges in K . So, the degree 4 vertices in $\text{lk}(3)$ are 1, 2 and 3. Then 1356, 1367, 2356, 2357, 3457 and 3467 are facets. Since $\text{lk}(7) \not\cong R_4$ and $\deg(78) = 4$, either $\text{lk}(7) \cong R_3$ or $\text{lk}(7)$ is an S_7^2 . In the former case, 2567 is a facet. This is not possible from $\text{lk}(25)$. So, $\text{lk}(7)$ is an S_7^2 . Then, from $\text{lk}(7)$, 1467, 2457 $\in K$. Now, from $\text{lk}(1)$, 1256 $\in K$. Here, $K = N_7$.

Case 2. Exactly one vertex whose link is isomorphic to R_4 and there exists a vertex whose link is isomorphic to R_3 . Using the same method as in Case 1, we find that $K \cong N_8$.

Case 3. Exactly one vertex whose link is isomorphic to R_4 , there is no vertex whose link is isomorphic to R_3 and there exists (at least) a vertex whose link is isomorphic to R_2 . Using the same method as in Case 1, we find that $K \cong N_9$.

Case 4. There is no vertex whose link is isomorphic to R_4 and there exist (at least) two vertices whose links are isomorphic to R_3 . Assume that $\text{lk}_K(8) = R_4$, so that $\deg(78) = 4$. Using the same method as in Case 1, we get the following: (i) if $\text{lk}_K(7) \cong R_3$ then $K = N_6$ and (ii) if $\text{lk}_K(7) \not\cong R_3$ then K is isomorphic to N_{10} or N_{11} .

Case 5. There is no vertex whose link is isomorphic to R_4 , there exists exactly one vertex whose link is isomorphic to R_3 and there exists (at least) a vertex whose link is isomorphic to R_2 . Using the same method as in Case 1, we find that K is isomorphic to N_{12} or N_{13} .

Case 6. There is no vertex whose link is isomorphic to R_4 or R_3 and there exist (at least) two vertices whose links are isomorphic to R_2 . Using the same method as in Case 1, we find that K is isomorphic to N_{14} or N_{15} . This completes the proof. $\square$

Proof of Theorem 2. Since $S_{8,m}^3$'s are combinatorial 3-manifolds and N_n 's are not combinatorial 3-manifolds, $S_{8,m}^3 \not\cong N_n$ for $35 \leq m \leq 38$, $1 \leq n \leq 15$. Part (a) now follows from Lemmas 3.1, 3.4. Part (b) follows from Lemmas 4.4, 4.5 and 4.6. $\square$

Lemma 4.7. Let $\mathcal{S}_0, \dots, \mathcal{S}_6$ be as in the proof of Lemma 3.2. If a combinatorial 3-manifold K is obtained from a member of $\mathcal{S}_j$ by a bistellar 2-move then K is isomorphic to a member of $\mathcal{S}_{j+1}$ for $0 \leq j \leq 5$. Moreover, no bistellar 2-move is possible from a member of $\mathcal{S}_6$.

Proof. Recall that $\mathcal{S}_0 = \{S_{8,35}^3, S_{8,36}^3, S_{8,37}^3, S_{8,38}^3\}$. The removable edges in $S_{8,37}^3$ are 13, 16, 17, 24, 27, 35, 46, 48 and 58. Since $(1, 4)(2, 7)(3, 8) \in \text{Aut}(S_{8,37}^3)$, up to isomorphisms it is

sufficient to consider the bistellar 2-moves κ_{27} , κ_{24} , κ_{48} , κ_{58} and κ_{46} only. Here $S_{8,33}^3 := \kappa_{27}(S_{8,37}^3)$, $S_{8,30}^3 := \kappa_{24}(S_{8,37}^3)$, $S_{8,32}^3 := \kappa_{48}(S_{8,37}^3)$, $S_{8,31}^3 := \kappa_{58}(S_{8,37}^3)$ and $\kappa_{46}(S_{8,37}^3) \cong S_{8,31}^3$ by the map $(1, 4, 5)(2, 7)(3, 6, 8)$.

The removable edges in $S_{8,38}^3$ are 13, 38, 78, 27, 25, 15 and 46. Since $(1, 2, 8)(7, 3, 5)$, $(1, 2)(3, 7)(4, 6) \in \text{Aut}(S_{8,38}^3)$, it is sufficient to consider the bistellar 2-moves κ_{46} and κ_{78} only. Here $S_{8,39}^3 := \kappa_{46}(S_{8,36}^3)$ and $\kappa_{78}(S_{8,38}^3) \cong S_{8,32}^3$ by the map $(1, 7, 8, 4, 6)(2, 3)$.

The removable edges in $S_{8,36}^3$ are 13, 35, 58, 68, 46, 24, 27, 17. Since $(1, 5, 6, 2)(3, 8, 4, 7)$ is an automorphism of $S_{8,36}^3$, it is sufficient to consider the bistellar 2-moves κ_{58} and κ_{68} only. Here $\kappa_{58}(S_{8,36}^3) = S_{8,31}^3$ and $\kappa_{68}(S_{8,36}^3) \cong S_{8,30}^3$ by the map $(1, 6, 4, 8, 2, 5, 7, 3)$.

The removable edges in $S_{8,35}^3$ are 13, 35, 57, 71, 24, 46, 68 and 82. Since $(1, 2, \dots, 8)$, $(1, 8)(2, 7)(3, 6)(4, 5) \in \text{Aut}(S_{8,35}^3)$, it is sufficient to consider the bistellar 2-moves κ_{68} only. Here $\kappa_{68}(S_{8,35}^3) \cong S_{8,30}^3$ by the map $(1, 7, 3)(2, 8, 4, 5, 6)$. This proves the result for $j = 0$.

By the same arguments as in the case for $j = 0$, one proves for the cases for $1 \leq j \leq 5$. We summarize these cases in the diagram given below. Last part follows from the fact that none of $S_{8,1}^3$, $S_{8,3}^3$ or $S_{8,3}^3$ has any removable edges. $\square$




Diagram 1 : Hasse diagram of the poset of the 8-vertex combinatorial 3-manifolds (the partial order relation is as defined in Section 2)

Lemma 4.8. *Let $\mathcal{N}_0, \dots, \mathcal{N}_3$ be as in the proof of Lemma 3.5. If a 3-pseudomanifold K is obtained from a member of $\mathcal{N}_j$ by a bistellar 2-move then K is isomorphic to a member of $\mathcal{N}_{j+1}$ for $0 \leq j \leq 2$. Moreover, no bistellar 2-move is possible from a member of $\mathcal{N}_3$.*

Proof. Recall that $\mathcal{N}_0 = \{N_1, \dots, N_{15}\}$. Since there are no degree 3 edges in N_1 , N_2 , N_5 and N_6 , no bistellar 2-moves are possible from N_1 , N_5 , N_6 or N_2 . The degree 3 edges in N_3 (respectively, in N_4) are 14, 16, 17, 36, 67 (respectively, 13, 35, 57, 72, 24, 46, 61). But, none of these edges is removable. So, bistellar 2-moves are not possible from N_3 or N_4 .

The removable edges in N_7 are 12, 14, 24, 56, 57 and 67. Since $(1, 2)(6, 7)$, $(1, 2)(5, 6)$ and $(1, 5)(2, 6)(3, 8)(4, 7)$ are automorphisms of N_7 , it follows that up to isomorphisms, we only have to consider the bistellar 2-move κ_{67} . Here, $N_{16} = \kappa_{67}(N_7)$.

The removable edges in N_8 are 15, 17, 24, 56, 57 and 67. Since $(1, 6)(2, 4)$, $(1, 6)(5, 7)$, $(2, 4)(5, 7) \in \text{Aut}(N_8)$, we only consider the bistellar 2-moves κ_{24} , κ_{56} and κ_{57} . Here, $N_{17} = \kappa_{24}(N_8)$, $N_{18} = \kappa_{56}(N_8)$ and $N_{19} = \kappa_{57}(N_8)$.

The removable edges in N_9 are 12, 23, 24 and 67. Since $(1, 4)(6, 7) \in \text{Aut}(N_9)$, we consider only κ_{12} , κ_{23} and κ_{67} . Here, $N_{21} = \kappa_{12}(N_9)$, $N_{23} = \kappa_{23}(N_9)$ and $\kappa_{67}(N_9) = N_{16}$.

The removable edges in N_{10} are 12, 14, 24, 56, 57 and 67. Since $(1, 7)(2, 5)(3, 8)(4, 6)$, $(1, 4)(6, 7) \in \text{Aut}(N_{10})$, we consider the bistellar 2-moves κ_{56} and κ_{57} only. Here, $N_{20} = \kappa_{56}(N_{10})$ and $\kappa_{67}(N_{10}) = N_{16}$.

The removable edges of N_{11} are 14, 24, 56, 57 and 67. Since $(1, 2)(5, 6)(3, 8) \in \text{Aut}(N_{11})$, we only consider the bistellar 2-moves κ_{14} , κ_{56} and κ_{67} . Here, $N_{22} = \kappa_{14}(N_{11})$, $\kappa_{56}(N_{11}) = N_{20}$ and $\kappa_{67}(N_{11}) \cong N_{18}$ (by the map $(2, 4)(5, 7)$).

The removable edges in N_{12} are 12, 23, 45 and 57. Here, $\kappa_{12}(N_{12}) \cong N_{22}$ (by the map $(2, 4, 6)$), $\kappa_{23}(N_{12}) = N_{23}$, $\kappa_{45}(N_{12}) \cong N_{21}$ (by the map $(1, 6, 5, 2, 7, 4)(3, 8)$) and $\kappa_{57}(N_{12}) \cong N_{18}$ (by the map $(1, 6, 7, 4)$).

The removable edges in N_{13} are 12, 23, 24, 56, 57 and 67. Since $(1, 4)(6, 7) \in \text{Aut}(N_{13})$, we only consider κ_{12} , κ_{23} , κ_{57} and κ_{67} . Here, $\kappa_{12}(N_{13}) \cong N_{22}$ (by the map $(2, 7, 5, 4)$), $\kappa_{23}(N_{13}) = N_{23}$, $\kappa_{57}(N_{13}) \cong N_{18}$ (by the map $(1, 4)(6, 7)$) and $\kappa_{67}(N_{13}) = N_{16}$.

The removable edges in N_{14} are 38, 56, 57, 67. Since $(1, 2, 4)(5, 6, 7)(3, 8) \in \text{Aut}(N_{14})$, we only consider κ_{38} and κ_{57} . Here, $N_{24} = \kappa_{38}(N_{14})$ and $\kappa_{57}(N_{14}) = N_{19}$.

The removable edges in N_{15} are 15, 23, 24, 58. Since $(1, 7)(2, 5)(3, 8)(4, 6) \in \text{Aut}(N_{15})$, we only consider the bistellar 2-moves κ_{23} and κ_{24} . Here, $\kappa_{23}(N_{15}) = N_{23}$ and $\kappa_{24}(N_{15}) \cong N_{21}$ (by the map $(1, 6, 5, 7, 4)$). This proves the result for $j = 0$.

By the same arguments as in the case for $j = 0$, one proves for the cases for $j = 1, 2$. We summarize these cases in the diagram given below. Last part follows from the fact that, for $N_i \in \mathcal{N}_3$, N_i has no removable edge. $\square$




Diagram 2: Hasse diagram of the poset of all the 3-pseudomanifolds $N_7, \dots, N_{35}$

Proof of Corollary 3. Let $\mathcal{S}_0, \dots, \mathcal{S}_6$ be as in the proof of Lemma 3.2. Let M be an 8-vertex combinatorial 3-manifold. Then, by Theorem 1, there exist bistellar 1-moves $\kappa_{A_1}, \dots, \kappa_{A_m}$, for some $m \geq 0$, such that $M_1 := \kappa_{A_m}(\dots(\kappa_{A_1}(M)))$ is a neighbourly 8-vertex 3-pseudomanifold. Since bistellar moves send a combinatorial 3-manifold to a combinatorial 3-manifold, M_1 is a combinatorial 3-manifold. Then, by Theorem 2, $M_1 \in \mathcal{S}_0$. In other words, $M = \kappa_{e_1}(\dots(\kappa_{e_m}(M_1)))$, where $M_1 \in \mathcal{S}_0$ and $\kappa_{e_m} : M_1 \mapsto \kappa_{e_m}(M_1)$, $\kappa_{e_i} : \kappa_{e_{i+1}}(\dots(\kappa_{e_m}(M_1))) \mapsto \kappa_{e_i}(\dots(\kappa_{e_m}(M_1)))$, for $1 \leq i \leq m - 1$ are bistellar 2-moves.

Therefore, by Lemma 4.7, $M \in \mathcal{S}_0 \cup \dots \cup \mathcal{S}_6$. The result now follows from Lemma 3.2. $\square$

Proof of Corollary 4. Let $\mathcal{N}_0, \dots, \mathcal{N}_3$ be as in the proof of Lemma 3.5. Let M be an 8-vertex normal 3-pseudomanifold. Then, by Theorem 1, there exist bistellar 1-moves $\kappa_{A_1}, \dots, \kappa_{A_m}$, for some $m \geq 0$, such that $M_1 := \kappa_{A_m}(\dots(\kappa_{A_1}(M)))$ is a neighbourly 3-pseudomanifold. Since bistellar moves send a normal 3-pseudomanifold to a normal 3-pseudomanifold, M_1 is normal. Hence, by Theorem 2, $M_1 \in \mathcal{N}_0$. In other words, $M = \kappa_{e_1}(\dots(\kappa_{e_m}(M_1)))$, where $M_1 \in \mathcal{N}_0$ and $\kappa_{e_m} : M_1 \mapsto \kappa_{e_m}(M_1)$, $\kappa_{e_i} : \kappa_{e_{i+1}}(\dots(\kappa_{e_m}(M_1))) \mapsto \kappa_{e_i}(\dots(\kappa_{e_m}(M_1)))$, for $1 \leq i \leq m-1$ are bistellar 2-moves. Therefore, by Lemma 4.8, $M \in \mathcal{N}_0 \cup \mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$. The result now follows from Lemma 3.5. $\square$

Acknowledgement : The authors thank the anonymous referees for many useful comments which helped to improve the presentation of this paper. The first author was partially supported by DST (Grant: SR/S4/MS-272/05) and by UGC-SAP/DSA-IV.

References

- [1] A. Altshuler, Combinatorial 3-manifolds with few vertices, *J. Combin. Theory (A)* **16** (1974), 165–173.
- [2] A. Altshuler, 3-pseudomanifolds with preassigned links, *Trans. Amer. Math. Soc.* **241** (1978), 213–237.
- [3] B. Bagchi and B. Datta, A structure theorem for pseudomanifolds, *Discrete Math.* **168** (1998), 41–60.
- [4] B. Bagchi and B. Datta, Minimal triangulations of sphere bundles over the circle, *J. Combin. Theory (A)* **115** (2008), 737–752.
- [5] B. Bagchi and B. Datta, Uniqueness of Walkup’s 9-vertex 3-dimensional Klein bottle, *Discrete Math.* (in press).
- [6] D. Barnette, The triangulations of the 3-sphere with up to 8 vertices, *J. Combin. Theory (A)* **14** (1973), 37–52.
- [7] B. Datta, Two dimensional weak pseudomanifolds on seven vertices, *Bol. Soc. Mat. Mexicana* **5** (1999), 419–426.
- [8] A. Emch, Triple and multiple systems, their geometric configurations and groups, *Trans. Amer. Math. Soc.* **31** (1929), 25–42.
- [9] E. Gawrilow and M. Joswig, polymake. <http://www.math.tu-berlin.de/polymake>, 1997–2007, Version 2.3.
- [10] B. Grünbaum and V. P. Sreedharan, An enumeration of simplicial 4-polytopes with 8 vertices, *J. Combin. Theory* **2** (1967), 437–465.
- [11] W. Kühnel, Minimal triangulations of Kummer varieties, *Abh. Math. Sem. Univ. Hamburg* **57** (1986), 7–20.
- [12] W. Kühnel, Triangulations of manifolds with few vertices, *Advances in Differential Geometry and Topology* (F. Tricerri, ed.), World Scientific, Singapore, 1990, pp. 59–114.
- [13] W. Kühnel, Topological aspects of twofold triple systems, *Expo. Math.* **16** (1998), 289–332.
- [14] F. H. Lutz, Triangulated manifolds with few vertices and vertex-transitive group actions, Thesis (TU, Berlin), Shaker-Verlag, Aachen, 1999.
- [15] D. W. Walkup, The lower bound conjecture for 3- and 4-manifolds, *Acta Math.* **125** (1970), 75–107.